



FACT SHEET

BIOCONTROL OF COMMON GRAPEVINE INSECT PESTS: AUSTRALIAN GRAPEVINE MOTH

By Dr Mary Retallack, Retallack Viticulture Pty Ltd



FINDING THE BALANCE... NATURALLY!

Healthy and diverse populations of predatory arthropods (insects and spiders) and parasitoids (wasps and flies) can help prevent grapevine pests from reaching economically damaging thresholds.

Growers can support healthy predator populations by providing a habitat that provides food, shelter and alternative prey/hosts and minimise the use of pesticides that are toxic to natural enemies.

Biocontrol options for common Australian grapevine pests are explored in this series of fact sheets. For a broader discussion about functional biodiversity please see the [EcoVineyards best practice management guide on functional biodiversity in Australian vineyards](#) and to read the other fact sheets in this series please visit the EcoVineyards [knowledge hub](#).

FAMILY NOCTUIDAE

Phalaenoides glycinae, Australian grapevine moth

DESCRIPTION: The Australian grapevine moth is a moth belonging to the family Noctuidae. The larvae may cause feeding damage to developing bunches and/or defoliation of grapevine leaves if populations are left unchecked. It is reported if caterpillars are accidentally fermented during the winemaking process, the resulting wine may have an elevated cineol (eucalyptol) concentration if the caterpillars have been eating Eucalyptus leaves.

DISTINCTIVE FEATURES: Grapevine moth eggs are commonly laid on the underside of leaves and resemble spherical domes. They are slightly ribbed and have a diameter of about 0.3 mm. The caterpillars grow to about 40 mm. They have striking black, white, yellow, and orange markings with a light brown head capsule and a big red rump. Adult moths have tufts of orange hair projecting from their abdomen and the base of their legs, contrasting with the black and white markings of their wings and body. The wingspan is about 50 mm.




Figure 1. (a) *Phalaenoides glycinae*, grapevine moth egg, (b) larva, and (c) adult [Photos: Mary Retallack].

BREEDING CYCLE: Moths emerge from over-wintering pupae in early spring and lay eggs on stems and leaves. There are two to three annual generations with larvae first typically appearing on vines in October, and the second generation of moths appearing in December. In areas with warm to hot summers, a third generation may occur between late summer and autumn.

WHEN TO MONITOR: Monitor from late October onwards.

SUGGESTED ACTION THRESHOLDS: Growers are encouraged to develop their own action thresholds based on data collected from monitoring and damage assessments at harvest over several seasons to reduce the likelihood of defoliation (especially later in the season leading up to harvest).

Biocontrol options

Many natural enemies, including insectivorous birds, microbats, predatory arthropods, and parasitic wasps, attack grapevine moth. Because it causes damage to vine foliage very rapidly once it is established, this pest is unlikely to be controlled satisfactorily through the reactive use of commercially available biological control agents. However, a functional diversity of predators will help to reduce the likelihood of populations reaching damaging levels. Alternatively, the strategic use of *Bt* sprays coupled with a feeding attractant may provide effective control.

PARASITOIDS: Parasitoids, such as tachinid flies and parasitoid wasps, contribute towards biocontrol. *Euplectrus agaristae*, chalcid wasp, is a parasitoid of the Australian vine moth larvae (Bernard et al., 2006a) and is present in Coonawarra vineyards along with *Echthromorpha intricatoria*, cream-spotted ichneumon, and *Lissopimpla semipunctata*, orchid dube wasp (Thomson and Hoffman, 2006).

PREDATORY ARTHROPODS: Important predators include *Cermatulus nasalis*, glossy shield bug, and *Oechalia schellenbergii*, predatory shield bug (Cordingley, 1981), ladybird larvae and spiders.

***Bacillus thuringiensis (Bt)*:** Bt products can be used anytime throughout the growing season. The *Bt* bacterium is toxic to moth larvae and, once consumed, it results in the paralysis of the digestive tract and larvae starve to death.




Figure 2. (a) ladybird beetle larva feeding on the Australian grapevine moth [Photo: Mary Retallack], (b) *Cermatulus nasalis*, glossy shield bug, [Landcare Research CC-BY 4.0], (c) *Oechalia schellenbergii*, predatory shield bug, and (d) *Cymbacha* sp., crab spider [Photos: Mary Retallack].

FURTHER READING

For more information on natural enemies, please see [natural predators of vineyards insect pests booklet](#) and associated [articles](#) and [fact sheets](#) on the [EcoVineyards knowledge hub](#).

REFERENCES

- Bernard M., Semerato L., Carter V., Wratten S.D. (2006) Beneficial insects in vineyards: Parasitoids of LBAM and grapevine moth in south-east Australia. *The Australian and New Zealand Grapegrower and Winemaker* 513:21-28.
- Cordingley C.L. (1981) Factors causing mortality of the grape vine moth, *Phalaenoides glycine* Lewin (Lepidoptera: Agaristidae). *Journal of the Australian Entomological Society*:33 -38.
- Thomson L.J., Hoffman A.A. (2006) Integrated strategies and bioindicators for sustainable grape production in Australia, in: D. Pimentel (Ed.), *Encyclopaedia of pest management*, Marcel Dekker, New York.

Disclaimer

The information contained in this EcoVineyards fact sheet is provided for information purposes only. Wine Australia and Retallack Viticulture Pty Ltd give no representations or warranties in relation to the content of the fact sheet including without limitation that it is without error or is appropriate for any particular purpose. No person should act in reliance on the content of this fact sheet without first obtaining specific, independent professional advice having regard to their site(s). Wine Australia and Retallack Viticulture Pty Ltd accept no liability for any direct or indirect loss or damage of any nature suffered or incurred in reliance on the content of the fact sheet.

For more information about the National EcoVineyards Program please visit www.ecovineyards.com.au @EcoVineyards

© Retallack Viticulture Pty Ltd, 2024



PROGRAM PARTNERS



REGIONAL PARTNERS



MARGARET RIVER WINE



The National EcoVineyards Program is funded by Wine Australia with levies from Australia's grape growers and winemakers and matching funds from the Australian Government.

ACKNOWLEDGEMENT OF COUNTRY

EcoVineyards proudly acknowledge the Aboriginal and Torres Strait Islander Peoples, and their ongoing cultural and spiritual connection to this ancient land on which we work and live.

As the Traditional custodians we recognise their wealth of ecological knowledge and the importance of caring for Country.

We pay our respect to elders past and present and extend this respect to all Aboriginal and Torres Strait Islander Peoples.



